

## The blob complex

Scott Morrison  
<http://tqft.net/>  
 joint work with Kevin Walker

UC Berkeley / Miller Institute for Basic Research


Low-Dimensional Topology and Categorification,  
 Stony Brook University, June 21-25 2010

slides: <http://tqft.net/talks>

paper: <http://tqft.net/blobs>

## What is the blob complex?

The blob complex takes an  $n$ -manifold  $\mathcal{M}$  and an ' $n$ -category with strong duality'  $\mathcal{C}$  and produces a chain complex,  $\mathcal{B}_*(\mathcal{M}; \mathcal{C})$ .



## $n$ -categories

There are many definitions of  $n$ -categories!

For most of what follows, I'll draw 2-dimensional pictures and rely on your intuition for pivotal categories.

We have yet another definition: *topological  $n$ -categories*

- ▶ A vector space  $\mathcal{C}(B^n)$  for every  $n$ -ball  $B$ .
- ▶ An associative gluing map: with  $B = \bigcup_i B_i$ , balls glued together to form a ball,

$$\bigotimes \mathcal{C}(B_i) \rightarrow \mathcal{C}(B)$$


(the  $\otimes$  is fibered over 'boundary restriction' maps).

- ▶ ...

These are easy to check for geometric examples, hard to check for algebraic examples.

## Cellulations of manifolds

Consider  $\text{cell}(M)$ , the category of cellulations of a manifold  $M$ , with morphisms 'antirefinements'.



An  $n$ -category  $\mathcal{C}$  gives a functor from  $\text{cell}(M)$  to vector spaces.


**objects** send a cellulation to the product of  $\mathcal{C}$  on each top-cell, restricting to the subset where boundaries agree

**morphisms** send an antirefinement to the appropriate gluing map.

## Fields

A field on  $\mathcal{M}^n$  is a choice of cellulation and a choice of  $n$ -morphism for each top-cell (with matching boundaries).

Example ( $\mathcal{C} = \text{TL}_d$  the Temperley-Lieb category)



Given a field on a ball, we can evaluate it to a morphism using the gluing map. We call the kernel the *null fields*.

$$\text{ev} \left( \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} - \frac{1}{d} \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} \right) = 0$$

## Background: TQFT invariants

### Definition

We associate to an  $n$ -manifold  $\mathcal{M}$  the skein module

$$\mathcal{A}(\mathcal{M}) = \mathcal{F}(\mathcal{M}) / \ker \text{ev},$$

fields modulo fields which evaluate to zero inside some ball.

Equivalently,  $\mathcal{A}(\mathcal{M})$  is the colimit of  $\mathcal{C}$  along  $\text{cell}(\mathcal{M})$ .

$\mathcal{A}(Y^{n-1} \times [0, 1])$  is a 1-category, and when  $Y \subset \partial X$ ,  $\mathcal{A}(X)$  is a module over  $\mathcal{A}(Y \times [0, 1])$ .

### Theorem (Gluing formula)

When  $Y \sqcup Y^{op} \subset \partial X$ ,

$$\mathcal{A}(X \bigcup_Y \text{circle}) \cong \mathcal{A}(X) \otimes_{\mathcal{A}(Y \times [0,1])} \text{circle}$$

## Motivation: Khovanov homology as a 4d TQFT

### Theorem

Khovanov homology gives a 4-category:

3-morphisms tangles, with the usual 3 operations,

4-morphisms  $\text{Hom}_{Kh}(T_1, T_2) = Kh(T_1 \cup \bar{T}_2)$ , composition defined by saddle cobordisms

There is a corresponding 4-manifold invariant. Given  $L \subset \partial W^4$ , it associates a doubly-graded vector space  $\mathcal{A}(W, L; Kh)$ .

$$\mathcal{A}(B^4, L; Kh) \cong Kh(L)$$

## Computations are hard

This invariant is hard to compute, because the TQFT skein module construction breaks the exact triangle for resolving a crossing.

$$\begin{array}{ccc} Kh(\text{crossing}) & & \mathcal{A}(M, \text{crossing}) \\ \swarrow & \nearrow & \vdots \\ Kh(\text{resol 1}) & \longrightarrow & Kh(\text{resol 2}) \\ & & \vdots \\ & & \mathcal{A}(M, \text{resol 1}) \text{ --- } \mathcal{A}(M, \text{resol 2}) \end{array}$$

There is a spectral sequence converging to 0 relating the blob homologies for the triangle of resolutions.

### Conjecture

It may be possible to compute the skein module by first computing the entire blob homology.


## Definition of the blob complex, $k = 0, 1$

### Motivation

A *local* construction, such that when  $\mathcal{M}$  is a ball,  $\mathcal{B}_*(\mathcal{M}; \mathcal{C})$  is a resolution of  $\mathcal{A}(\mathcal{M}; \mathcal{C})$ .

$\mathcal{B}_0(\mathcal{M}; \mathcal{C}) = \mathcal{F}(\mathcal{M})$ , arbitrary fields on  $\mathcal{M}$ .

$$\mathcal{B}_1(\mathcal{M}; \mathcal{C}) = \mathbb{C} \left\{ (B, u, r) \mid \begin{array}{l} B \text{ an embedded ball} \\ u \in \mathcal{F}(B) \text{ in the kernel} \\ r \in \mathcal{F}(\mathcal{M} \setminus B) \end{array} \right\}.$$



$$d_1 : (B, u, r) \mapsto u \circ r \quad \mathcal{B}_0 / \text{im}(d_1) \cong \mathcal{A}(\mathcal{M}; \mathcal{C})$$

## Definition, $k = 2$

$$\mathcal{B}_2 = \mathcal{B}_2^{\text{disjoint}} \oplus \mathcal{B}_2^{\text{nested}}$$

$$\mathcal{B}_2^{\text{disjoint}} = \mathbb{C} \left\{ \left( \text{Diagram} \mid \text{ev}_{B_i}(u_i) = 0 \right) \right\}$$

$$d_2 : (B_1, B_2, u_1, u_2, r) \mapsto (B_2, u_2, r \circ u_1) - (B_1, u_1, r \circ u_2)$$

$$\mathcal{B}_2^{\text{nested}} = \mathbb{C} \left\{ \left( \text{Diagram} \mid \text{ev}_{B_1}(u) = 0 \right) \right\}$$

$$d_2 : (B_1, B_2, u, r', r) \mapsto (B_2, u \circ r', r) - (B_1, u, r \circ r')$$

## Definition, general case

$$\mathcal{B}_k = \mathbb{C} \left\{ \left( \text{Diagram} \right) \right\}$$

$k$  blobs, properly nested or disjoint, with “innermost” blobs labelled by fields that evaluate to zero.

$$d_k : \mathcal{B}_k \rightarrow \mathcal{B}_{k-1} = \sum_i (-1)^i (\text{erase blob } i)$$

## Hochschild homology

TQFT on  $S^1$  is ‘coinvariants’

$$\mathcal{A}(S^1, A) = \mathbb{C} \left\{ \left( \text{Diagram} \right) \right\} / \left\{ \left( \text{Diagram} \right) - \left( \text{Diagram} \right) \right\} = A / (ab - ba)$$

Blob homology on  $S^1$  is Hochschild homology

The Hochschild complex is ‘coinvariants of the bar resolution’

$$\dots \rightarrow A \otimes A \otimes A \rightarrow A \otimes A \xrightarrow{m \otimes a \rightarrow ma - am} A$$

We check universal properties, as it’s hard to directly construct an isomorphism.

## An action of $C_*(\text{Homeo}(\mathcal{M}))$

### Theorem

There's a chain map

$$C_*(\text{Homeo}(\mathcal{M})) \otimes \mathcal{B}_*(\mathcal{M}) \rightarrow \mathcal{B}_*(\mathcal{M}).$$

which is associative up to homotopy, and compatible with gluing.

Taking  $H_0$ , this is the mapping class group acting on a TQFT skein module.

$$H_0(\text{Homeo}(\mathcal{M})) \otimes \mathcal{A}(\mathcal{M}) \rightarrow \mathcal{A}(\mathcal{M}).$$

## Gluing

$\mathcal{B}_*(Y \times [0, 1])$  is naturally an  $A_\infty$  category

multiplication ( $m_2$ ): gluing  $[0, 1] \simeq [0, 1] \cup [0, 1]$

associativity up to homotopy ( $m_k$ ): reparametrising  $[0, 1]$  using the action of  $C_*(\text{Homeo}([0, 1]))$ .

If  $Y \subset \partial X$  then  $\mathcal{B}_*(X)$  is an  $A_\infty$  module over  $\mathcal{B}_*(Y)$ .

### Theorem (Gluing formula)

When  $Y \sqcup Y^{op} \subset \partial X$ ,

$$\mathcal{B}_*(X \bigcup_Y \circlearrowleft) \cong \mathcal{B}_*(X) \overset{A_\infty}{\otimes}_{\mathcal{B}_*(Y)} \circlearrowleft.$$

In principle, we can compute blob homology from a handle decomposition, by iterated Hochschild homology.

## Higher Deligne conjecture


### Deligne conjecture

Chains on the little discs operad acts on Hochschild cohomology.

Call  $\text{Hom}_{\mathcal{B}_*(\partial M)}(\mathcal{B}_*(\mathcal{M}), \mathcal{B}_*(\mathcal{M}))$  'blob cochains on  $\mathcal{M}$ '.

### Theorem (Higher Deligne conjecture)

Chains on the  $n$ -dimensional fat graph operad acts on blob cochains.



## Maps to a space

Fix a target space  $\mathcal{T}$ . There is an  $A_\infty$   $n$ -category  $\pi_{\leq n}^\infty(\mathcal{T})$  defined by

$$\pi_{\leq n}^\infty(\mathcal{T})(B) = C_*(\text{Maps}(B \rightarrow \mathcal{T})).$$

(Here  $B$  is an  $n$ -ball.)

### Theorem

The blob complex recovers mapping spaces:

$$\mathcal{B}_*(\mathcal{M}; \pi_{\leq n}^\infty(\mathcal{T})) \cong C_*(\text{Maps}(\mathcal{M} \rightarrow \mathcal{T}))$$

This generalizes a result of Lurie: if  $\mathcal{T}$  is  $n - 1$  connected,  $\pi_{\leq n}^\infty(\mathcal{T})$  is an  $E_n$ -algebra and in this special case the blob complex is presumably the same as his topological chiral homology.