




# The Blob Complex, part 2

Kevin Walker

(joint work with Scott Morrison)



slides and preprint available at [canyon23.net/math/](http://canyon23.net/math/)

(or the URLs Scott gave)

## Goals:

- n-category definition optimized for TQFTs
- should be very easy to show that topological examples satisfy the axioms
- as simple as possible (but not simpler)
- both plain and infinity type categories
- also define modules, coends, tensor products, etc.

## Goals:

- n-category definition optimized for TQFTs
- should be very easy to show that topological examples satisfy the axioms
- as simple as possible (but not simpler)
- both plain and infinity type categories
- also define modules, coends, tensor products, etc.

## Main ideas:

- don't skeletonize (don't try to minimize generators, don't try to minimize relations)
- build in "strong" duality from the start
- non-recursive (don't need to know what an  $(n-1)$ -category is)




## Ingredients for an n-category:

1. morphisms in dimensions 0 through n
2. domain/range/boundary
3. composition
4. identity morphisms
5. special behavior in dimension n

# Morphisms


- Need to decide on “shape” of morphisms





# Morphisms

- Need to decide on “shape” of morphisms



- We will allow morphisms to be of **any** shape, so long as it is homeomorphic to a ball

**Morphisms (preliminary version):** *For any  $k$ -manifold  $X$  homeomorphic to the standard  $k$ -ball, we have a set of  $k$ -morphisms  $\mathcal{C}_k(X)$ .*



**Morphisms (preliminary version):** *For any  $k$ -manifold  $X$  homeomorphic to the standard  $k$ -ball, we have a set of  $k$ -morphisms  $\mathcal{C}_k(X)$ .*

**Morphisms:** *For each  $0 \leq k \leq n$ , we have a functor  $\mathcal{C}_k$  from the category of  $k$ -balls and homeomorphisms to the category of sets and bijections.*

**Morphisms (preliminary version):** *For any  $k$ -manifold  $X$  homeomorphic to the standard  $k$ -ball, we have a set of  $k$ -morphisms  $\mathcal{C}_k(X)$ .*

**Morphisms:** *For each  $0 \leq k \leq n$ , we have a functor  $\mathcal{C}_k$  from the category of  $k$ -balls and homeomorphisms to the category of sets and bijections.*

Balls could be PL, topological, or smooth. Also unoriented, oriented, Spin, Pin $_{\pm}$ . We will concentrate on the case of PL unoriented balls.



# Examples

Let  $T$  be a topological space.

$\mathcal{C}_k(X^k) = \text{Maps}(X \rightarrow T)$ , for  $k < n$ ,  $X$  a  $k$ -ball.

$\mathcal{C}_n(X^n) = \text{Maps}(X \rightarrow T)$  modulo homotopy rel boundary  
(fundamental  $n$ -groupoid of  $T$ )

$\mathcal{C}_k(X^k) = \text{Maps}(X \rightarrow T)$ , for  $k < n$ ,  $X$  a  $k$ -ball.

$\mathcal{C}_n(X^n) = C_*(\text{Maps}(X \rightarrow T))$  (singular chains)  
( $\infty$  version of fundamental groupoid of  $T$ )

$\mathcal{C}_k(X^k) = \{\text{embedded decorated cell complexes in } X\}$ , for  $k < n$ .

$\mathcal{C}_n(X^n) = \{\text{embedded decorated cell complexes in } X\}$  modulo isotopy and other local relations

More examples

$$\bigcirc = q^5 + q^4 + q + 1 + q^{-1} + q^{-4} + q^{-5}$$

$$\bigcirc\bigcirc = q^9 + q^6 + q^5 + q^4 + q^3 + q + 2 + q^{-1} + q^{-3} + q^{-4} + q^{-5} + q^{-6} + q^{-9}$$

$$\text{loop} = 0$$

$$\text{figure-eight} = -(q^3 + q^2 + q + q^{-1} + q^{-2} + q^{-3}) \text{---}$$

$$\text{triangle} = (q^2 + 1 + q^{-2}) \text{---}$$

$$\text{square} = -(q + q^{-1}) \left( \text{---} + \text{---} \right) + (q + 1 + q^{-1}) \left( \text{---} + \text{---} \right) \left( \text{---} \right)$$

$$\text{pentagon} = - \left( \text{---} + \text{---} + \text{---} + \text{---} + \text{---} \right) + \left( \text{---} + \text{---} + \text{---} + \text{---} + \text{---} \right)$$

$$\text{cross} = \left( \text{---} - \text{---} - \frac{1}{q^2 - 1 + q^{-2}} \right) \left( + \frac{1}{q + 1 + q^{-1}} \text{---} \right)$$

(Kuperberg)




# More examples

Let  $A$  be a traditional linear  $n$ -category with strong duality (e.g. pivotal 2-category).

$\mathcal{C}_k(X^k) = \{A\text{-string diagrams in } X\}$ , for  $k < n$ .

$\mathcal{C}_n(X^n) = \{\text{finite linear combinations of } A\text{-string diagrams in } X\}$  modulo diagrams which evaluate to zero



$\mathcal{C}_k(X^k) = \{A\text{-string diagrams in } X\}$ , for  $k < n$ .

$\mathcal{C}_n(X^n) = \text{blob complex of } X \text{ based on } A\text{-string diagrams}$

**Boundaries (domain and range), part 1:** *For each  $0 \leq k \leq n - 1$ , we have a functor  $\mathcal{C}_k$  from the category of  $k$ -spheres and homeomorphisms to the category of sets and bijections.*



**Boundaries (domain and range), part 1:** *For each  $0 \leq k \leq n - 1$ , we have a functor  $\mathcal{C}_k$  from the category of  $k$ -spheres and homeomorphisms to the category of sets and bijections.*

**Boundaries, part 2:** *For each  $k$ -ball  $X$ , we have a map of sets  $\partial : \mathcal{C}(X) \rightarrow \mathcal{C}(\partial X)$ . These maps, for various  $X$ , comprise a natural transformation of functors.*


**Boundaries (domain and range), part 1:** For each  $0 \leq k \leq n - 1$ , we have a functor  $\mathcal{C}_k$  from the category of  $k$ -spheres and homeomorphisms to the category of sets and bijections.

**Boundaries, part 2:** For each  $k$ -ball  $X$ , we have a map of sets  $\partial : \mathcal{C}(X) \rightarrow \mathcal{C}(\partial X)$ . These maps, for various  $X$ , comprise a natural transformation of functors.

**Domain + range  $\rightarrow$  boundary:** Let  $S = B_1 \cup_E B_2$ , where  $S$  is a  $k$ -sphere ( $0 \leq k \leq n - 1$ ),  $B_i$  is a  $k$ -ball, and  $E = B_1 \cap B_2$  is a  $k-1$ -sphere. Let  $\mathcal{C}(B_1) \times_{\mathcal{C}(E)} \mathcal{C}(B_2)$  denote the fibered product of the two maps  $\partial : \mathcal{C}(B_i) \rightarrow \mathcal{C}(E)$ . Then (axiom) we have an injective map

$$\text{gl}_E : \mathcal{C}(B_1) \times_{\mathcal{C}(E)} \mathcal{C}(B_2) \rightarrow \mathcal{C}(S)$$

which is natural with respect to the actions of homeomorphisms.




- Let  $\mathcal{C}(S)_E \subset \mathcal{C}(S)$  denote the image of  $\text{gl}_E$



- Let  $\mathcal{C}(S)_E \subset \mathcal{C}(S)$  denote the image of  $\text{gl}_E$ 
  - Given  $c \in \mathcal{C}(\partial(X))$ , let  $\mathcal{C}(X; c) \stackrel{\text{def}}{=} \partial^{-1}(c)$

- Let  $\mathcal{C}(S)_E \subset \mathcal{C}(S)$  denote the image of  $\text{gl}_E$
- Given  $c \in \mathcal{C}(\partial(X))$ , let  $\mathcal{C}(X; c) \stackrel{\text{def}}{=} \partial^{-1}(c)$
- Given  $E \subset \partial X$ , let  $\mathcal{C}(X)_E \stackrel{\text{def}}{=} \partial^{-1}(\mathcal{C}(\partial X)_E)$




- Let  $\mathcal{C}(S)_E \subset \mathcal{C}(S)$  denote the image of  $\text{gl}_E$

- Given  $c \in \mathcal{C}(\partial(X))$ , let  $\mathcal{C}(X; c) \stackrel{\text{def}}{=} \partial^{-1}(c)$

- Given  $E \subset \partial X$ , let  $\mathcal{C}(X)_E \stackrel{\text{def}}{=} \partial^{-1}(\mathcal{C}(\partial X)_E)$

- In most examples, we require that the sets  $\mathcal{C}(X; c)$  (for all  $n$ -balls  $X$  and all boundary conditions  $c$ ) have extra structure, e.g. vector space or chain complex






**Composition:** Let  $B = B_1 \cup_Y B_2$ , where  $B$ ,  $B_1$  and  $B_2$  are  $k$ -balls ( $0 \leq k \leq n$ ) and  $Y = B_1 \cap B_2$  is a  $k-1$ -ball. Let  $E = \partial Y$ , which is a  $k-2$ -sphere. Note that each of  $B$ ,  $B_1$  and  $B_2$  has its boundary split into two  $k-1$ -balls by  $E$ . We have restriction (domain or range) maps  $\mathcal{C}(B_i)_E \rightarrow \mathcal{C}(Y)$ . Let  $\mathcal{C}(B_1)_E \times_{\mathcal{C}(Y)} \mathcal{C}(B_2)_E$  denote the fibered product of these two maps. Then (axiom) we have a map

$$\text{gl}_Y : \mathcal{C}(B_1)_E \times_{\mathcal{C}(Y)} \mathcal{C}(B_2)_E \rightarrow \mathcal{C}(B)_E$$



which is natural with respect to the actions of homeomorphisms, and also compatible with restrictions to the intersection of the boundaries of  $B$  and  $B_i$ . If  $k < n$  we require that  $\text{gl}_Y$  is injective. (For  $k = n$ , see below.)



**Composition:** Let  $B = B_1 \cup_Y B_2$ , where  $B$ ,  $B_1$  and  $B_2$  are  $k$ -balls ( $0 \leq k \leq n$ ) and  $Y = B_1 \cap B_2$  is a  $k-1$ -ball. Let  $E = \partial Y$ , which is a  $k-2$ -sphere. Note that each of  $B$ ,  $B_1$  and  $B_2$  has its boundary split into two  $k-1$ -balls by  $E$ . We have restriction (domain or range) maps  $\mathcal{C}(B_i)_E \rightarrow \mathcal{C}(Y)$ . Let  $\mathcal{C}(B_1)_E \times_{\mathcal{C}(Y)} \mathcal{C}(B_2)_E$  denote the fibered product of these two maps. Then (axiom) we have a map

$$\text{gl}_Y : \mathcal{C}(B_1)_E \times_{\mathcal{C}(Y)} \mathcal{C}(B_2)_E \rightarrow \mathcal{C}(B)_E$$


which is natural with respect to the actions of homeomorphisms, and also compatible with restrictions to the intersection of the boundaries of  $B$  and  $B_i$ . If  $k < n$  we require that  $\text{gl}_Y$  is injective. (For  $k = n$ , see below.)



**Strict associativity:** The composition (gluing) maps above are strictly associative.



**Multi-composition:** Given any decomposition  $B = B_1 \cup \dots \cup B_m$  of a  $k$ -ball into small  $k$ -balls, there is a map from an appropriate subset (like a fibered product) of  $\mathcal{C}(B_1) \times \dots \times \mathcal{C}(B_m)$  to  $\mathcal{C}(B)$ , and these various  $m$ -fold composition maps satisfy an operad-type strict associativity condition.





**Product (identity) morphisms:** Let  $X$  be a  $k$ -ball and  $D$  be an  $m$ -ball, with  $k + m \leq n$ . Then we have a map  $\mathcal{C}(X) \rightarrow \mathcal{C}(X \times D)$ , usually denoted  $a \mapsto a \times D$  for  $a \in \mathcal{C}(X)$ . If  $f : X \rightarrow X'$  and  $\tilde{f} : X \times D \rightarrow X' \times D'$  are maps such that the diagram

$$\begin{array}{ccc} X \times D & \xrightarrow{\tilde{f}} & X' \times D' \\ \pi \downarrow & & \downarrow \pi \\ X & \xrightarrow{f} & X' \end{array}$$

commutes, then we have

$$\tilde{f}(a \times D) = f(a) \times D'.$$

Product morphisms are compatible with gluing (composition) in both factors:

$$(a' \times D) \bullet (a'' \times D) = (a' \bullet a'') \times D$$

and

$$(a \times D') \bullet (a \times D'') = a \times (D' \bullet D'').$$

Product morphisms are associative:




$$(a \times D) \times D' = a \times (D \times D').$$

(Here we are implicitly using functoriality and the obvious homeomorphism  $(X \times D) \times D' \rightarrow X \times (D \times D')$ .) Product morphisms are compatible with restriction:


$$\text{res}_{X \times E}(a \times D) = a \times E$$

for  $E \subset \partial D$  and  $a \in \mathcal{C}(X)$ .


We need something a little more general  
than plain products




We need something a little more general than plain products



We need something a little more general than plain products



“extended isotopy”





## Plain n-cat:

**Extended isotopy invariance in dimension  $n$ :** *Let  $X$  be an  $n$ -ball and  $f : X \rightarrow X$  be a homeomorphism which restricts to the identity on  $\partial X$  and is extended isotopic (rel boundary) to the identity. Then  $f$  acts trivially on  $\mathcal{C}(X)$ .*

## Plain n-cat:

**Extended isotopy invariance in dimension  $n$ :** *Let  $X$  be an  $n$ -ball and  $f : X \rightarrow X$  be a homeomorphism which restricts to the identity on  $\partial X$  and is extended isotopic (rel boundary) to the identity. Then  $f$  acts trivially on  $\mathcal{C}(X)$ .*


## Infinity n-cat:

**Families of homeomorphisms act in dimension  $n$ .** *For each  $n$ -ball  $X$  and each  $c \in \mathcal{C}(\partial X)$  we have a map of chain complexes*

$$C_*(\text{Homeo}_{\partial}(X)) \otimes \mathcal{C}(X; c) \rightarrow \mathcal{C}(X; c).$$

*Here  $C_*$  means singular chains and  $\text{Homeo}_{\partial}(X)$  is the space of homeomorphisms of  $X$  which fix  $\partial X$ . These action maps are required to be associative up to homotopy, and also compatible with composition (gluing).*

# Equivalences between this n-cat definition and more traditional ones (at least for $n=1$ or $2$ )




# Colimit construction

- Let  $\mathcal{C}$  be in  $n$ -category.
- We want to extend  $\mathcal{C}$  to arbitrary  $k$ -manifolds  $Y$ ,  $0 \leq k \leq n$ .




# Colimit construction

- Let  $\mathcal{C}$  be in  $n$ -category.
- We want to extend  $\mathcal{C}$  to arbitrary  $k$ -manifolds  $Y$ ,  $0 \leq k \leq n$ .
- Let  $\mathcal{J}$  be the category (partially ordered set) whose objects are decompositions of  $Y$  into balls and whose morphisms are anti-refinements (coarsenings) of these decompositions.





# Colimit construction



- Let  $\mathcal{C}$  be in  $n$ -category.
- We want to extend  $\mathcal{C}$  to arbitrary  $k$ -manifolds  $Y$ ,  $0 \leq k \leq n$ .
- Let  $\mathcal{J}$  be the category (partially ordered set) whose objects are decompositions of  $Y$  into balls and whose morphisms are anti-refinements (coarsenings) of these decompositions.
- There is a functor which assigns to a decomposition  $Y = \bigcup_i X_i$  the set (or vector space or chain complex)  $\bigotimes_i \mathcal{C}(X_i)$ .

# Colimit construction

- Let  $\mathcal{C}$  be in  $n$ -category.
- We want to extend  $\mathcal{C}$  to arbitrary  $k$ -manifolds  $Y$ ,  $0 \leq k \leq n$ .
- Let  $\mathcal{J}$  be the category (partially ordered set) whose objects are decompositions of  $Y$  into balls and whose morphisms are anti-refinements (coarsenings) of these decompositions.
- There is functor which assigns to a decomposition  $Y = \bigcup_i X_i$  the set (or vector space or chain complex)  $\bigotimes_i \mathcal{C}(X_i)$ .
  - Define  $\mathcal{C}(Y)$  to be the colimit (or homotopy colimit) of this functor.



# Newfangled blob complex

- Given an  $A_\infty$   $n$ -category  $\mathcal{C}$ , we define the newfangled blob complex of an  $n$ -manifold  $M$  to be the above homotopy colimit  $\mathcal{C}(M)$ .



# Newfangled blob complex

- Given an  $A_\infty$   $n$ -category  $\mathcal{C}$ , we define the newfangled blob complex of an  $n$ -manifold  $M$  to be the above homotopy colimit  $\mathcal{C}(M)$ .
- Given a plain  $n$ -category  $\mathcal{C}$ , we can construct an  $A_\infty$   $n$ -category  $\mathcal{D}$  by defining  $\mathcal{D}(X) = \mathcal{B}_*^{\mathcal{C}}(X)$  for each  $n$ -ball  $X$ .

# Newfangled blob complex

- Given an  $A_\infty$   $n$ -category  $\mathcal{C}$ , we define the newfangled blob complex of an  $n$ -manifold  $M$  to be the above homotopy colimit  $\mathcal{C}(M)$ .
- Given a plain  $n$ -category  $C$ , we can construct an  $A_\infty$   $n$ -category  $\mathcal{D}$  by defining  $\mathcal{D}(X) = \mathcal{B}_*^C(X)$  for each  $n$ -ball  $X$ .
- $\mathcal{D}$  is in some sense the free resolution of  $C$  as an  $A_\infty$   $n$ -category.

# Newfangled blob complex

- Given an  $A_\infty$   $n$ -category  $\mathcal{C}$ , we define the newfangled blob complex of an  $n$ -manifold  $M$  to be the above homotopy colimit  $\mathcal{C}(M)$ .
- Given a plain  $n$ -category  $C$ , we can construct an  $A_\infty$   $n$ -category  $\mathcal{D}$  by defining  $\mathcal{D}(X) = \mathcal{B}_*^C(X)$  for each  $n$ -ball  $X$ .
- $\mathcal{D}$  is in some sense the free resolution of  $C$  as an  $A_\infty$   $n$ -category.
  - Let  $M^n = F^{n-k} \times Y^k$ . Let  $C$  be a plain  $n$ -category. Let  $\mathcal{F}$  be the  $A_\infty$   $k$ -category which assigns to a  $k$ -ball  $X$  the old-fashioned blob complex  $\mathcal{B}_*^C(X \times F)$ .

# Newfangled blob complex

- Given an  $A_\infty$   $n$ -category  $\mathcal{C}$ , we define the newfangled blob complex of an  $n$ -manifold  $M$  to be the above homotopy colimit  $\mathcal{C}(M)$ .
- Given a plain  $n$ -category  $C$ , we can construct an  $A_\infty$   $n$ -category  $\mathcal{D}$  by defining  $\mathcal{D}(X) = \mathcal{B}_*^C(X)$  for each  $n$ -ball  $X$ .
- $\mathcal{D}$  is in some sense the free resolution of  $C$  as an  $A_\infty$   $n$ -category.
  - Let  $M^n = F^{n-k} \times Y^k$ . Let  $C$  be a plain  $n$ -category. Let  $\mathcal{F}$  be the  $A_\infty$   $k$ -category which assigns to a  $k$ -ball  $X$  the old-fashioned blob complex  $\mathcal{B}_*^C(X \times F)$ .
  - Theorem:  $\mathcal{F}(Y) \simeq \mathcal{B}_*^C(F \times Y)$ .



# Newfangled blob complex


- Given an  $A_\infty$   $n$ -category  $\mathcal{C}$ , we define the newfangled blob complex of an  $n$ -manifold  $M$  to be the above homotopy colimit  $\mathcal{C}(M)$ .
- Given a plain  $n$ -category  $C$ , we can construct an  $A_\infty$   $n$ -category  $\mathcal{D}$  by defining  $\mathcal{D}(X) = \mathcal{B}_*^C(X)$  for each  $n$ -ball  $X$ .
- $\mathcal{D}$  is in some sense the free resolution of  $C$  as an  $A_\infty$   $n$ -category.
  - Let  $M^n = F^{n-k} \times Y^k$ . Let  $C$  be a plain  $n$ -category. Let  $\mathcal{F}$  be the  $A_\infty$   $k$ -category which assigns to a  $k$ -ball  $X$  the old-fashioned blob complex  $\mathcal{B}_*^C(X \times F)$ .
  - Theorem:  $\mathcal{F}(Y) \simeq \mathcal{B}_*^C(F \times Y)$ .
- Corollary:  $\mathcal{D}(M) \simeq \mathcal{B}_*^C(M)$  for any  $n$ -manifold  $M$ . (Proof: Let  $F$  above be a point.) So the old-fashioned and newfangled blob complexes are homotopy equivalent.

# Modules

- Let  $\mathcal{C}$  be an  $n$ -category.
- Modules for  $\mathcal{C}$  are defined in a similar style.


# Modules

- Let  $\mathcal{C}$  be an  $n$ -category.
- Modules for  $\mathcal{C}$  are defined in a similar style.
- A *marked  $k$ -ball* is a pair  $(B, M)$  which is homeomorphic to the standard pair  $(B^k, B^{k-1})$ .



# Modules

- Let  $\mathcal{C}$  be an  $n$ -category.
- Modules for  $\mathcal{C}$  are defined in a similar style.
- A *marked  $k$ -ball* is a pair  $(B, M)$  which is homeomorphic to the standard pair  $(B^k, B^{k-1})$ .




- A  $\mathcal{C}$ -module  $\mathcal{M}$  is a collection of functors  $\mathcal{M}_k$  from the category of marked  $k$ -balls to the category of sets,  $0 \leq k \leq n$ .



# Modules

- Let  $\mathcal{C}$  be an  $n$ -category.
- Modules for  $\mathcal{C}$  are defined in a similar style.
- A *marked  $k$ -ball* is a pair  $(B, M)$  which is homeomorphic to the standard pair  $(B^k, B^{k-1})$ .




- A  $\mathcal{C}$ -module  $\mathcal{M}$  is a collection of functors  $\mathcal{M}_k$  from the category of marked  $k$ -balls to the category of sets,  $0 \leq k \leq n$ .
- In the top dimension  $n$  we have the same extra structure as  $\mathcal{C}$  (vector space, chain complex, ...).


- Motivating example: Let  $W$  be an  $m+1$ -manifold with non-empty boundary. Let  $\mathcal{E}$  be an  $m+n$ -category.
- Let  $\mathcal{C}$  be the  $n$ -category with  $\mathcal{C}(X) \stackrel{\text{def}}{=} \mathcal{E}(X \times \partial W)$ .

- Motivating example: Let  $W$  be an  $m+1$ -manifold with non-empty boundary. Let  $\mathcal{E}$  be an  $m+n$ -category.
- Let  $\mathcal{C}$  be the  $n$ -category with  $\mathcal{C}(X) \stackrel{\text{def}}{=} \mathcal{E}(X \times \partial W)$ .
  - Define the  $\mathcal{C}$ -module  $\mathcal{M}$  by


$$\mathcal{M}(M, B) \stackrel{\text{def}}{=} \mathcal{E} \left( (B \times \partial W) \cup_{M \times \partial W} (M \times W) \right).$$



- Two different ways of cutting a marked  $k$ -ball into two pieces, so two different kinds of composition. (One is composition within  $\mathcal{M}$ , the other is the action of  $\mathcal{C}$  on  $\mathcal{M}$ .)




action




$\mathcal{M}$ -composition



- Two different ways of cutting a marked  $k$ -ball into two pieces, so two different kinds of composition. (One is composition within  $\mathcal{M}$ , the other is the action of  $\mathcal{C}$  on  $\mathcal{M}$ .)





action




$\mathcal{M}$ -composition


- Various kinds of mixed strict associativity.



- Two different ways of cutting a marked  $k$ -ball into two pieces, so two different kinds of composition. (One is composition within  $\mathcal{M}$ , the other is the action of  $\mathcal{C}$  on  $\mathcal{M}$ .)





action



$\mathcal{M}$ -composition


- Various kinds of mixed strict associativity.




- $\mathcal{M}$  can be thought of as a collection of  $n-1$ -categories with some extra structure.



- Two different ways of cutting a marked  $k$ -ball into two pieces, so two different kinds of composition. (One is composition within  $\mathcal{M}$ , the other is the action of  $\mathcal{C}$  on  $\mathcal{M}$ .)





action



$\mathcal{M}$ -composition

- Various kinds of mixed strict associativity.



- $\mathcal{M}$  can be thought of as a collection of  $n-1$ -categories with some extra structure.
- For  $n = 1, 2$  this is equivalent to the usual notion of module.

# Decorated colimit construction

- Let  $W$  be a  $k$ -manifold. Let  $Y_i$  be a collection of disjoint codimension 0 submanifolds of  $\partial W$ .
- Let  $\mathcal{C}$  be an  $n$ -category and  $\mathcal{N} = \{\mathcal{N}_i\}$  be a collection of  $\mathcal{C}$ -modules, thought of as labels of  $\{Y_i\}$ .




# Decorated colimit construction

- Let  $W$  be a  $k$ -manifold. Let  $Y_i$  be a collection of disjoint codimension 0 submanifolds of  $\partial W$ .
- Let  $\mathcal{C}$  be an  $n$ -category and  $\mathcal{N} = \{\mathcal{N}_i\}$  be a collection of  $\mathcal{C}$ -modules, thought of as labels of  $\{Y_i\}$ .
- We can use a variation on the above colimit construction to define a set (or vector space or chain complex if  $k = n$ )  $\mathcal{C}(W, \mathcal{N})$ .

# Decorated colimit construction


- Let  $W$  be a  $k$ -manifold. Let  $Y_i$  be a collection of disjoint codimension 0 submanifolds of  $\partial W$ .
- Let  $\mathcal{C}$  be an  $n$ -category and  $\mathcal{N} = \{\mathcal{N}_i\}$  be a collection of  $\mathcal{C}$ -modules, thought of as labels of  $\{Y_i\}$ .
- We can use a variation on the above colimit construction to define a set (or vector space or chain complex if  $k = n$ )  $\mathcal{C}(W, \mathcal{N})$ .
- The object of the colimit are decompositions of  $W$  into (plain) balls  $X_j$  and marked balls  $(B_l, M_l)$ , with  $M_l = B_l \cap \{Y_i\}$ .





# Decorated colimit construction


- Let  $W$  be a  $k$ -manifold. Let  $Y_i$  be a collection of disjoint codimension 0 submanifolds of  $\partial W$ .
- Let  $\mathcal{C}$  be an  $n$ -category and  $\mathcal{N} = \{\mathcal{N}_i\}$  be a collection of  $\mathcal{C}$ -modules, thought of as labels of  $\{Y_i\}$ .
- We can use a variation on the above colimit construction to define a set (or vector space or chain complex if  $k = n$ )  $\mathcal{C}(W, \mathcal{N})$ .
- The object of the colimit are decompositions of  $W$  into (plain) balls  $X_j$  and marked balls  $(B_l, M_l)$ , with  $M_l = B_l \cap \{Y_i\}$ .



- This defines an  $n-k$ -category which assigns  $\mathcal{C}(D \times W, \mathcal{N})$  to a ball  $D$ . (Here  $\mathcal{N}_i$  labels  $D \times Y_i$ .)

# Tensor products and gluing

- As a simple special case of this construction, given  $\mathcal{C}$ -modules  $\mathcal{N}_1$  and  $\mathcal{N}_2$ , define the tensor product  $\mathcal{N}_1 \otimes \mathcal{N}_2$  (an  $n-1$ -category) to be the result of taking  $W$  to be an interval and letting  $\mathcal{N}_1$  and  $\mathcal{N}_2$  label the endpoints of the interval.





# Tensor products and gluing

- As a simple special case of this construction, given  $\mathcal{C}$ -modules  $\mathcal{N}_1$  and  $\mathcal{N}_2$ , define the tensor product  $\mathcal{N}_1 \otimes \mathcal{N}_2$  (an  $n-1$ -category) to be the result of taking  $W$  to be an interval and letting  $\mathcal{N}_1$  and  $\mathcal{N}_2$  label the endpoints of the interval.



- Gluing theorem: Let  $M^{n-k} = M_1 \cup_Y M_2$ . Let  $\mathcal{C}$  be an  $n$ -category. The above constructions give a  $k$ -category  $\mathcal{C}(M)$ , a  $k-1$ -category  $\mathcal{C}(Y)$ , and two  $\mathcal{C}(Y)$ -modules  $\mathcal{C}(M_i)$ . Then

$$\mathcal{C}(M) \simeq \mathcal{C}(M_1) \otimes_{\mathcal{C}(Y)} \mathcal{C}(M_2).$$

