

Math 3325, 2013

Problem Set 2

Discuss in tutorial on August 5 and 12

These questions are all about the problem of finding the closest point in  $L^p([0, 1])$ , in the subspace

$$S_p = \{f \in L^p([0, 1]) \mid \int_0^1 xf(x) dx = 0\},$$

to the function  $g(x) = 1$ .

1. Show that  $S_p$  is a closed subspace of  $L^p([0, 1])$ , for all  $p \in [1, \infty)$ .

2. For  $p = 2$ , find the closest point in  $S_2$  to  $g$  (in the  $L^2$  metric).

3. For  $p = 1$ , show that for every  $\epsilon > 0$  there is an  $f_n \in S_1$  such that  $\|g - f_n\|_1 \leq 1/2 + \epsilon$ , but that there is no  $f \in S_1$  such that  $\|g - f\|_1 \leq 1/2$ . In particular, there is no closest point to  $g$  in  $S_1$  in the  $L^1$  metric.

4. What happens in  $L^p$  for  $1 < p < 2$ ? Hint: if  $g = f + h$ , where  $f \in S_p$ , use Hölder's inequality

$$\left| \int_0^1 u(x)v(x) dx \right| \leq \|u\|_{L^p([0,1])} \|v\|_{L^q([0,1])}, \quad p^{-1} + q^{-1} = 1,$$

to get a lower bound on  $\|h\|_p$ . Then use the fact that equality in Hölder's inequality will occur if  $u, v \geq 0$  and  $u = cv^{q-1}$ .