

Overview

In preparation for the exam, we'll look at the questions asked on the 2013 Mid-Semester Exam.

Sample Question: Lines & Planes

Let P be the plane in \mathbb{R}^3 defined by the equation $2x + y - z = 1$, and let L be the line through the point $(1, 1, 1)$ which is orthogonal to P .

- 1 Find an equation for P of the form $\mathbf{n} \cdot (\mathbf{r} - \mathbf{r}_0) = 0$ for some vector \mathbf{n} and some vector \mathbf{r}_0 .
- 2 Find an equation for L .
- 3 Let Q be the plane containing L and the point $(1, 1, 2)$. Find an equation for Q .

Solution: Lines & Planes

Let P be the plane in \mathbb{R}^3 defined by the equation $2x + y - z = 1$, and let L be the line through the point $(1, 1, 1)$ which is orthogonal to P .

- 1 Find an equation for P of the form $\mathbf{n} \cdot (\mathbf{r} - \mathbf{r}_0) = 0$ for some vector \mathbf{n} and some vector \mathbf{r}_0 .

To find the equation of a plane P , we need a **normal vector** to P and a **point** on P .

The plane $Ax + By + Cz + D = 0$ has normal vector $\begin{bmatrix} A \\ B \\ C \end{bmatrix}$, so a normal

vector to P is given by $\begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix}$. To find a point on P , we can plug in

$x = y = 0$ and see that $(0, 0, -1)$ satisfies the equation $2x + y - z = 1$.

Thus the general formula $\mathbf{n} \cdot (\mathbf{r} - \mathbf{r}_0) = 0$ becomes

$$\begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z + 1 \end{bmatrix} = 0.$$

Solution: Lines & Planes

Let P be the plane in \mathbb{R}^3 defined by the equation $2x + y - z = 1$, and let L be the line through the point $(1, 1, 1)$ which is orthogonal to P .

- ② Find an equation for L .

A direction vector for L is any normal vector to P : i.e., any scalar multiple

of $\mathbf{n} = \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix}$. This yields the vector equation

$$\mathbf{r} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + t \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix},$$

with the associated parametric equations

$$x = 1 + 2t \quad y = 1 + t \quad z = 1 - t.$$

Solution: Lines & Planes

Let P be the plane in \mathbb{R}^3 defined by the equation $2x + y - z = 1$, and let L be the line through the point $(1, 1, 1)$ which is orthogonal to P .

- ③ Let Q be the plane containing L and the point $(1, 1, 2)$. Find an equation for Q .

To find a normal vector to the new plane, take the cross product of two vectors parallel to Q . For example, you could choose a direction vector for

L and the vector $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ between the two given points on Q :

$$\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 1 & -1 \\ 0 & 0 & 1 \end{vmatrix} = \mathbf{i} - 2\mathbf{j}.$$

Any equation for the plane is acceptable, including the following:

$$\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix} - \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} \right) \cdot \begin{bmatrix} 1 \\ -2 \\ 0 \end{bmatrix} = 0,$$

Sample Question: Bases & Coordinates

The set $\mathcal{B} = \{t + 1, 1 + t^2, 3 - t^2\}$ is a basis for \mathbb{P}_2 .

- ① If $p(t) = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}_{\mathcal{B}}$, express p in the form $p(t) = a + bt + ct^2$.
- ② Find the coordinate vector of the polynomial $q(t) = 2 - 2t$ with respect to \mathcal{B} coordinates.

Solution: Bases & Coordinates

The set $\mathcal{B} = \{t + 1, 1 + t^2, 3 - t^2\}$ is a basis for \mathbb{P}_2 .

- 1 If $p(t) = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}_{\mathcal{B}}$, express p in the form $p(t) = a + bt + ct^2$.

Since the \mathcal{B} coordinates of p are 1, 1, and -1 , we have

$$p(t) = 1(t + 1) + 1(1 + t^2) - 1(3 - t^2) = -1 + t + 2t^2.$$

Solution: Bases & Coordinates

The set $\mathcal{B} = \{t + 1, 1 + t^2, 3 - t^2\}$ is a basis for \mathbb{P}_2 .

- 2 Find the coordinate vector of the polynomial $q(t) = 2 - 2t$ with respect to \mathcal{B} coordinates.

We need a, b , and c such that

$$a(t + 1) + b(1 + t^2) + c(3 - t^2) = 2 - 2t.$$

Collecting like powers of t gives us a system of equations:

$$\begin{aligned} a + b + 3c &= 2 \\ a &= -2 \\ b - c &= 0. \end{aligned}$$

The unique solution to this is $a = -2, b = c = 1$.

To protect against algebra mistakes, check that

$$-2(t + 1) + 1(1 + t^2) + 1(3 - t^2) = 2 - 2t.$$

Sample Question: Vector Spaces

Decide whether each of the following sets is a vector space. If it is a vector space, state its dimension. If it is not a vector space, explain why.

- 1 A is the set of 2×2 matrices whose entries are integers.
- 2 B is the set of vectors in \mathbb{R}^3 which are orthogonal to $\begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$.
- 3 C is the set of polynomials whose derivative is 0:

$$C = \{p(x) \in \mathbb{P} \mid \frac{d}{dx}p(x) = 0\}.$$

Solution: Vector Spaces

Decide whether each of the following sets is a vector space. If it is a vector space, state its dimension. If it is not a vector space, explain why.

- ① A is the set of 2×2 matrices whose entries are integers.

This is a subset of the vector space of 2×2 matrices with real entries, so we can check if the three subspace axioms hold:

- ① Is $\mathbf{0}$ in the set?
- ② Is the set closed under addition?
- ③ Is the set closed under scalar multiplication?

No, this is not a vector space. This set is not closed under multiplication by a non-integer scalar. For example,

$$\frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & 0 \end{bmatrix} \text{ is not in } A.$$

Solution: Vector Spaces

Decide whether each of the following sets is a vector space. If it is a vector space, state its dimension. If it is not a vector space, explain why.

- ② B is the set of vectors in \mathbb{R}^3 which are orthogonal to $\begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$.

As before, we could check the 3 subspace axioms, but it's quicker to observe that B is the null space of the matrix $[1 \ 0 \ 2]$, and the null space of a matrix is always a subspace.

We can find a basis for the null space explicitly and check that it has 2 vectors. Alternatively, observe that the matrix $[1 \ 0 \ 2]$ has rank 1, so its null space is two-dimensional by the Rank Theorem.

Checking the 3 subspace axioms

① $\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = 0$, so $\mathbf{0} \in B$.

② Suppose $\mathbf{v}, \mathbf{u} \in B$. Then $\mathbf{v} \cdot \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = \mathbf{u} \cdot \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = 0$.

$$(\mathbf{u} + \mathbf{v}) \cdot \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = \mathbf{u} \cdot \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} + \mathbf{v} \cdot \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = 0 + 0 = 0.$$

Since $\mathbf{u} + \mathbf{v}$ is in B , B is closed under addition.

- ③ Suppose $\mathbf{v} \in B$.

$$(c\mathbf{v}) \cdot \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = c \left(\mathbf{v} \cdot \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} \right) = c \cdot 0 = 0.$$

Since $c\mathbf{v}$ is in B , B is closed under scalar multiplication.

Solution: Vector Spaces

Decide whether each of the following sets is a vector space. If it is a vector space, state its dimension. If it is not a vector space, explain why.

- the set of polynomials whose derivative is 0:

$$C = \left\{ p(x) \in \mathbb{P} \mid \frac{d}{dx} p(x) = 0 \right\}.$$

We can solve this problem by recognising that the polynomials whose derivatives are 0 are exactly the constant polynomials, so $C = \mathbb{R}^1$. It follows that C is a one-dimensional vector space.

It is also acceptable to show that C is a subspace of the vector space \mathbb{P} by verifying each of the subspace axioms.

Sample Question: Linear transformations

A linear transformation $T : M_{2 \times 2} \rightarrow M_{2 \times 2}$ is defined by:

$$T \left(\begin{bmatrix} a & b \\ c & d \end{bmatrix} \right) = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}.$$

(a) Calculate $T \left(\begin{bmatrix} a & b \\ c & d \end{bmatrix} \right)$.

- (b) Which, if any, of the following matrices are in $\ker(T)$?

$$\begin{bmatrix} 1 & 1 \\ 3 & 3 \end{bmatrix} \quad \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} \quad \begin{bmatrix} 1 & 3 \\ 1 & 3 \end{bmatrix}$$

- (c) Which, if any, of the following matrices are in $\text{range}(T)$?

$$\begin{bmatrix} -2 & 2 \\ 2 & -2 \end{bmatrix} \quad \begin{bmatrix} 1 & -1 \\ -2 & 2 \end{bmatrix} \quad \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

- (d) Find the kernel of T and explain why T is not one to one.

- (e) Explain why T does not map $M_{2 \times 2}$ onto $M_{2 \times 2}$.

Sample Question: Subspaces associated to a matrix

Consider the matrix A :

$$\begin{bmatrix} 2 & -4 & 0 & 2 \\ -1 & 2 & 1 & 2 \\ 1 & -2 & 1 & 4 \end{bmatrix}.$$

- (i) Find a basis for $\text{Nul } A$.
- (ii) Find a basis for $\text{Col } A$.
- (iii) Consider the linear transformation $T_A : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ defined by $T_A(\mathbf{x}) = A\mathbf{x}$. Give a geometric description of the range of T_A as a subspace of \mathbb{R}^3 . What is its dimension? Does it pass through the origin?

We begin by row-reducing A :

$$\begin{bmatrix} 2 & -4 & 0 & 2 \\ -1 & 2 & 1 & 2 \\ 1 & -2 & 1 & 4 \end{bmatrix} \xrightarrow{\text{rref}} \begin{bmatrix} 1 & -2 & 0 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

(i) Find a basis for $\text{Nul } A$.

The general solution to $R \begin{bmatrix} w \\ x \\ y \\ z \end{bmatrix} = 0$ is $y + 3z = 0$, $w - 2x + z = 0$, so

$$\text{Nul } A = \left\{ \begin{bmatrix} 2x - z \\ x \\ -3z \\ z \end{bmatrix} \right\} = \left\{ x \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + z \begin{bmatrix} -1 \\ 0 \\ -3 \\ 1 \end{bmatrix} \right\}$$

and so $\mathcal{B} = \left\{ \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ -3 \\ 1 \end{bmatrix} \right\}$ is a basis for $\text{Nul } A$.

We begin by row-reducing A :

$$\begin{bmatrix} 2 & -4 & 0 & 2 \\ -1 & 2 & 1 & 2 \\ 1 & -2 & 1 & 4 \end{bmatrix} \xrightarrow{\text{rref}} \begin{bmatrix} 1 & -2 & 0 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

(ii) Find a basis for $\text{Col } A$.

A basis for $\text{Col } A$ is obtained by taking every column of A that corresponds to a pivot column in the row reduced form of A . Thus the first and third columns

$$\mathcal{C} = \left\{ \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \right\}$$

form a basis for $\text{Col } A$.

(iii) Consider the linear transformation $T_A : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ defined by $T_A(\mathbf{x}) = A\mathbf{x}$. Give a geometric description of the range of T_A as a subspace of \mathbb{R}^3 . What is its dimension? Does it pass through the origin?

The range of T_A is exactly the column space of A . We just saw that it has a basis with two elements, so it is two dimensional. It is a plane in \mathbb{R}^3 , and passed through the origin, because every vector subspace contains $\mathbf{0}$.

Revision: Definitions

- What is a vector space? Give some examples.
- What is a subspace? How do you check if a subset of a vector space is a subspace?
- What is a linear transformation? Give some examples.
- What does it mean for a set of vectors to be linearly independent? How do you check this?
- What are the coordinates of a vector with respect to a basis?

Revision: Geometry of \mathbb{R}^3

- What information do you need to determine a line? A plane?
- How can you check if two lines are orthogonal? Parallel?
- How do you find the distance between a point and a line? A point and a plane?
- How can you find the angle between two vectors?
- What are the scalar and vector projections of one vector onto another? Can you describe these in words?

Revision: Bases

- What is a basis for a vector space?
- If the dimension of V is n , then V and \mathbb{R}^n are *isomorphic*. What does this mean and how do we know it's true?
- In an n -dimensional vector space,
 - ▶ any n linearly independent vectors form a basis.
 - ▶ any n vectors which span V form a basis.
 - ▶ any set of vectors which spans V contains a basis for V .
 - ▶ any set of linearly independent vectors can be extended to a basis for V .
- How do you find a basis for the null space of a matrix? The column space? The row space? The kernel of the associated linear transformation? (Which pair of these are the same?)