

## Overview

Last week we introduced vectors in Euclidean space and the operations of vector addition, scalar multiplication, dot product, and (for  $\mathbb{R}^3$ ) cross product.

### Question

*How can we use vectors to describe lines and planes in  $\mathbb{R}^3$ ?*

(From Stewart §10.5)

## Warm-up

### Question

*Describe all the vectors in  $\mathbb{R}^3$  which are orthogonal to the 0 vector. Can you rephrase your answer as a statement about solutions to some linear equation?*

Remember that the statement " $\mathbf{v}$  is orthogonal to  $\mathbf{u}$ " is equivalent to " $\mathbf{v} \cdot \mathbf{u} = 0$ ".

This question asks for all the vectors  $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$  such that  $\begin{bmatrix} x \\ y \\ z \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = 0$ .

Using the definition of the dot product, this translates to asking what

$\begin{bmatrix} x \\ y \\ z \end{bmatrix}$  satisfy the equation  $0x + 0y + 0z = 0$ ...

...the answer is that all vectors in  $\mathbb{R}^3$  are orthogonal to the 0 vector.

Equivalently, every triple  $(x, y, z)$  is a solution to the linear equation  $0x + 0y + 0z = 0$ .

## Lines in $\mathbb{R}^2$

In the  $xy$ -plane the general form of the equation of a line is

$$ax + by = c,$$

where  $a$  and  $b$  are not both zero. If  $b \neq 0$  then this equation can be rewritten as

$$y = -(a/b)x + c/b,$$

which has the form  $y = mx + k$ . (Here  $m$  is the slope of the line and the point  $(0, k)$  is its  $y$ -intercept.)

### Example 1

Let  $L$  be the line  $2x + y = 3$ . The line has slope  $m = -2$  and the  $y$ -intercept is  $(0, 3)$ .

Alternatively, we could think about this line ( $y = -2x + 3$ ) as the path traced out by a moving particle.

Suppose that the particle is initially at the point  $(0, 3)$  at time  $t = 0$ . Suppose, too, that its  $x$ -coordinate changes at a constant rate of 1 unit per second and its  $y$ -coordinate changes as a constant rate of  $-2$  units per second.

At  $t = 1$  the particle is at  $(1, 1)$ . If we assume it's always been moving this way, then we also know that at  $t = -2$  it was at  $(-2, 7)$ . In general, we can display the relationship in vector form:

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} t \\ -2t + 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \end{bmatrix} + t \begin{bmatrix} 1 \\ -2 \end{bmatrix}$$

What is the significance of the vector  $\mathbf{v} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$ ?

In this expression,  $\mathbf{v}$  is a vector parallel to the line  $L$ , and is called a *direction vector* for  $L$ . The previous example shows that we can express  $L$  in terms of a direction vector and a vector to specific point on  $L$ :

#### Definition

The equation

$$\mathbf{r} = \mathbf{r}_0 + t\mathbf{v}$$

is the *vector equation* of the line  $L$ . The variable  $t$  is called a *parameter*.

Here,  $\mathbf{r}_0$  is the vector to a specific point on  $L$ ; any vector  $\mathbf{r}$  which satisfies this equation is a vector to some point on  $L$ .

#### Example 2

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \end{bmatrix} + t \begin{bmatrix} 1 \\ -2 \end{bmatrix} \quad (1)$$

is the *vector equation* of the line  $L$ .

If we express the vectors in a vector equation for  $L$  in components, we get a collection of equations relating scalars.

#### Definition

For  $\mathbf{r} = \begin{bmatrix} x \\ y \end{bmatrix}$ ,  $\mathbf{r}_0 = \begin{bmatrix} x_0 \\ y_0 \end{bmatrix}$ ,  $\mathbf{v} = \begin{bmatrix} a \\ b \end{bmatrix}$ , the *parametric equations* of the line  $\mathbf{r} = \mathbf{r}_0 + t\mathbf{v}$  are

$$\begin{aligned} x &= x_0 + ta \\ y &= y_0 + tb. \end{aligned}$$

## Lines in $\mathbb{R}^3$

The definitions of the vector and parametric forms of a line carry over perfectly to  $\mathbb{R}^3$ .

### Definition

The *vector form of the equation of the line  $L$*  in  $\mathbb{R}^2$  or  $\mathbb{R}^3$  is

$$\mathbf{r} = \mathbf{r}_0 + t\mathbf{v}$$

where  $\mathbf{r}_0$  is a specific point on  $L$  and  $\mathbf{v} \neq \mathbf{0}$  is a direction vector for  $L$ . The equations corresponding to the components of the vector form of the equation are called *parametric equations* of  $L$ .

### Example 3

Let  $\mathbf{r}_0 = \begin{bmatrix} 1 \\ 4 \\ -2 \end{bmatrix}$  and  $\mathbf{v} = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$ . Then the vector equation of the line  $L$  is

$$\mathbf{r} = \begin{bmatrix} 1 \\ 4 \\ -2 \end{bmatrix} + t \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}.$$

The line  $L$  contains the point  $(1, 4, -2)$  and has direction parallel to

$\mathbf{v} = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$ . By taking different values of  $t$  we can find different points on the line.

### Question

*For a given line, is the vector equation for the line unique?*

No, any vector parallel to the direction vector is another direction vector, and each choice of a point on  $L$  will give a different  $\mathbf{r}_0$ .

#### Example 4

The line with parametric equations

$$x = 1 + 2t \quad y = -4t \quad z = -3 + 5t.$$

can also be expressed as

$$x = 3 + 2t \quad y = -4 - 4t \quad z = 2 + 5t.$$

or as

$$x = 1 - 4t \quad y = 8t \quad z = -3 - 10t.$$

Note that a fixed value of  $t$  corresponds to three different points on  $L$  when plugged into the three different systems.

### Symmetric equations of a line

Another way of describing a line  $L$  is to eliminate the parameter  $t$  from the parametric equations

$$x = x_0 + at \quad y = y_0 + bt \quad z = z_0 + ct$$

If  $a \neq 0$ ,  $b \neq 0$  and  $c \neq 0$  then we can solve each of the scalar equations for  $t$  and obtain

$$\frac{x - x_0}{a} = \frac{y - y_0}{b} = \frac{z - z_0}{c}.$$

These equations are called the *symmetric equations* of the line  $L$  through  $(x_0, y_0, z_0)$  parallel to  $\mathbf{v}$ . The numbers  $a, b$  and  $c$  are called the *direction numbers* of  $L$ .

If, for example  $a = 0$ , the equation becomes

$$x = x_0, \quad \frac{y - y_0}{b} = \frac{z - z_0}{c}.$$

#### Example 5

Find parametric and symmetric equations for the line through  $(1, 2, 3)$  and parallel to  $2\mathbf{i} + 3\mathbf{j} - 4\mathbf{k}$ .

The line has the vector parametric form

$$\mathbf{r} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k} + t(2\mathbf{i} + 3\mathbf{j} - 4\mathbf{k}),$$

or scalar parametric equations

$$\begin{cases} x = 1 + 2t \\ y = 2 + 3t \\ z = 3 - 4t \end{cases} \quad (-\infty < t < \infty).$$

Its symmetric equations are

$$\frac{x - 1}{2} = \frac{y - 2}{3} = \frac{z - 3}{-4}.$$

### Example 6

Determine whether the two lines given by the parametric equations below intersect

$$L_1 : x = 1 + 2t, y = 3t, z = 2 - t$$

$$L_2 : x = -1 + s, y = 4 + s, z = 1 + 3s$$

If  $L_1$  and  $L_2$  intersect, there will be values of  $s$  and  $t$  satisfying

$$1 + 2t = -1 + s$$

$$3t = 4 + s$$

$$2 - t = 1 + 3s$$


Solving the first two equations gives  $s = 14, t = 6$ , but these values don't satisfy the third equation. We conclude that the lines  $L_1$  and  $L_2$  don't intersect.

In fact, their direction vectors are not proportional, so the lines aren't parallel, either. They are *skew* lines.


### Planes in $\mathbb{R}^3$

We described a line as the set of position vectors expressible as  $\mathbf{r}_0 + \mathbf{v}$ , where  $\mathbf{r}_0$  was a position vector of a point in  $L$  and  $\mathbf{v}$  was any vector parallel to  $L$ .

We can describe a plane the same way: the set of position vectors expressible as the sum of a position vector to a point in  $P$  and an arbitrary vector parallel to  $P$ .



Choose a vector  $\mathbf{n}$  which is orthogonal to the plane and choose an arbitrary point  $P_0$  in the plane.



How can we use this data to describe all the other points  $P$  which lie in the plane?

Let  $\mathbf{r}_0$  and  $\mathbf{r}$  be the position vectors of  $P_0$  and  $P$  respectively.

The normal vector  $\mathbf{n}$  is orthogonal to every vector in the plane. In particular  $\mathbf{n}$  is orthogonal to  $\mathbf{r} - \mathbf{r}_0$  and so we have

$$\mathbf{n} \cdot (\mathbf{r} - \mathbf{r}_0) = 0.$$

This equation

$$\mathbf{n} \cdot (\mathbf{r} - \mathbf{r}_0) = 0. \quad (2)$$

can be rewritten as

$$\mathbf{n} \cdot \mathbf{r} = \mathbf{n} \cdot \mathbf{r}_0. \quad (3)$$

Either of the equations (2) or (3) is called a *vector equation of the plane*.

### Example 7

Find a vector equation for the plane passing through  $P_0 = (0, -2, 3)$  and normal to the vector  $\mathbf{n} = 4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}$ .

We have  $\mathbf{r}_0 = \langle 0, -2, 3 \rangle$  and  $\mathbf{n} = \langle 4, 2, -3 \rangle$ . Thus the vector form is

$$\mathbf{n} \cdot (\mathbf{r} - \mathbf{r}_0) = 0,$$

or

$$(4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}) \cdot [(x - 0)\mathbf{i} + (y + 2)\mathbf{j} + (z - 3)\mathbf{k}] = 0.$$

Expanding this gives us a *scalar equation* for the plane...

Given  $\mathbf{n} = \langle A, B, C \rangle$ ,  $\mathbf{r} = \langle x, y, z \rangle$  and  $\mathbf{r}_0 = \langle x_0, y_0, z_0 \rangle$ , the vector equation  $\mathbf{n} \cdot (\mathbf{r} - \mathbf{r}_0) = 0$  becomes

$$\langle A, B, C \rangle \cdot \langle x - x_0, y - y_0, z - z_0 \rangle = 0,$$

or

$$A(x - x_0) + B(y - y_0) + C(z - z_0) = 0. \quad (4)$$

Equation (4) is the *scalar equation of the plane* through  $P_0(x_0, y_0, z_0)$  with normal vector  $\mathbf{n} = \langle A, B, C \rangle$ .

The equation

$$A(x - x_0) + B(y - y_0) + C(z - z_0) = 0.$$

can be written more simply in **standard form**

$$Ax + By + Cz + D = 0,$$

where  $D = -(Ax_0 + By_0 + Cz_0)$ .

If  $D = 0$ , the plane passes through the origin.

### Example 8

Find a scalar equation for the plane passing through  $P_0 = (0, -2, 3)$  and normal to the vector  $\mathbf{n} = 4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}$ .

The vector form is

$$(4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}) \cdot [(x - 0)\mathbf{i} + (y + 2)\mathbf{j} + (z - 3)\mathbf{k}] = 0,$$

which in scalar form becomes

$$4(x - 0) + 2(y + 2) - 3(z - 3) = 0$$

and this is equivalent to

$$4x + 2y - 3z = -13.$$

### Example 9

Find a scalar equation of the plane containing the points

$$P = (1, 1, 2), \quad Q = (0, 2, 3), \quad R = (-1, -1, -4).$$

First, we should find a normal vector  $\mathbf{n}$  to the plane, and there are several ways to do this.

The vector  $\mathbf{n} = n_1\mathbf{i} + n_2\mathbf{j} + n_3\mathbf{k}$  will be perpendicular to  $\overrightarrow{PQ} = -\mathbf{i} + \mathbf{j} + \mathbf{k}$  and  $\overrightarrow{PR} = -2\mathbf{i} - 2\mathbf{j} - 6\mathbf{k}$ . Therefore, we can solve a system of linear equations:

$$0 = \mathbf{n} \cdot (-\mathbf{i} + \mathbf{j} + \mathbf{k}) = -n_1 + n_2 + n_3$$

$$0 = \mathbf{n} \cdot (-2\mathbf{i} - 2\mathbf{j} - 6\mathbf{k}) = -2n_1 - 2n_2 - 6n_3.$$

One solution to this system is  $\mathbf{n} = -\mathbf{i} - 2\mathbf{j} + \mathbf{k}$ , so this is an example of a normal vector to the plane containing the 3 given points.

We can use this normal vector  $\mathbf{n} = -\mathbf{i} - 2\mathbf{j} + \mathbf{k}$ , together with any one of the given points to write the equation of the plane. Using  $Q = \begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}$ , the equation is

$$-(x - 0) - 2(y - 2) + 1(z - 3) = 0,$$

which simplifies to

$$x + 2y - z = 1.$$

The first step in this example was finding the normal vector  $\mathbf{n}$ , but in fact, there's another way to do this.

Recall that *in*  $\mathbb{R}^3$  *only*, there is a product of two vectors called a *cross product*. The cross product of  $\mathbf{a}$  and  $\mathbf{b}$  is a vector denoted  $\mathbf{a} \times \mathbf{b}$  which is orthogonal to both  $\mathbf{a}$  and  $\mathbf{b}$ . If we have two nonzero vectors  $\mathbf{a}$  and  $\mathbf{b}$  parallel to our plane, then  $\mathbf{n} = \mathbf{a} \times \mathbf{b}$  is a normal vector.

### Example 10

Consider the two planes

$$x - y + z = -1 \quad \text{and} \quad 2x + y + 3z = 4.$$

Explain why the planes above are not parallel and find a direction vector for the line of intersection.

Two planes are parallel if and only if their normal vectors are parallel. Normal vectors for the two planes above are for example

$$\mathbf{n}_1 = \mathbf{i} - \mathbf{j} + \mathbf{k} \quad \text{and} \quad \mathbf{n}_2 = 2\mathbf{i} + \mathbf{j} + 3\mathbf{k}$$

respectively. These vectors are not parallel, so the planes can't be parallel and must intersect. A vector  $\mathbf{v}$  parallel to the line of intersection is a vector which is orthogonal to both the normal vectors above. We can find such a vector by calculating the cross product of the normal vectors:

$$\mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -1 & 1 \\ 2 & 1 & 3 \end{vmatrix} = -4\mathbf{i} - \mathbf{j} + 3\mathbf{k}.$$



### Example 11

Find the line through the origin and parallel to the line of intersection of the two planes

$$x + 2y - z = 2 \quad \text{and} \quad 2x - y + 4z = 5.$$

The planes have respective normals

$$\mathbf{n}_1 = \mathbf{i} + 2\mathbf{j} - \mathbf{k} \quad \text{and} \quad \mathbf{n}_2 = 2\mathbf{i} - \mathbf{j} + 4\mathbf{k}.$$

A direction vector for their line of intersection is given by

$$\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = 7\mathbf{i} - 6\mathbf{j} - 5\mathbf{k}.$$

A vector parametric equation of the line is

$$\mathbf{r} = t(7\mathbf{i} - 6\mathbf{j} - 5\mathbf{k}),$$

since the line passes through the origin.

Parametric equations for this line are, for example,

$$\begin{aligned}x &= 7t \\y &= -6t \\z &= -5t\end{aligned}$$

and the corresponding symmetric equations are

$$\frac{x}{7} = \frac{y}{-6} = \frac{z}{-5}.$$

### Recommended exercises for review

Stewart §10.5: 1, 3, 15, 19, 25, 29, 35