

## Eigenvectors and eigenvalues

From Lay, §5.1

### Overview

Most of the material we've discussed so far falls loosely under two headings:

- geometry of  $\mathbb{R}^n$ , and
- generalisation of 1013 material to abstract vector spaces.

Today we'll begin our study of eigenvectors and eigenvalues. This is fundamentally different from material you've seen before, but we'll draw on the earlier material to help us understand this central concept in linear algebra. This is also one of the topics that you're most likely to see applied in other contexts.

### Question

*If you want to understand a linear transformation, what's the smallest amount of information that tells you something meaningful?*

This is a very vague question, but studying eigenvalues and eigenvectors gives us one way to answer it.

From Lay, §5.1

### Definition

An *eigenvector* of an  $n \times n$  matrix  $A$  is a non-zero vector  $\mathbf{x}$  such that  $A\mathbf{x} = \lambda\mathbf{x}$  for some scalar  $\lambda$ .

An *eigenvalue* of an  $n \times n$  matrix  $A$  is a scalar  $\lambda$  such that  $A\mathbf{x} = \lambda\mathbf{x}$  has a non-zero solution; such a vector  $\mathbf{x}$  is called an *eigenvector corresponding to*  $\lambda$ .

### Example 1

$$\text{Let } A = \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix}.$$

Then any nonzero vector  $\begin{bmatrix} x \\ 0 \end{bmatrix}$  is an eigenvector for the eigenvalue 3:

$$\begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ 0 \end{bmatrix} = \begin{bmatrix} 3x \\ 0 \end{bmatrix}.$$

Similarly, any nonzero vector  $\begin{bmatrix} 0 \\ y \end{bmatrix}$  is an eigenvector for the eigenvalue 2.

Sometimes it's not as obvious what the eigenvectors are.

### Example 2

$$\text{Let } B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}.$$

Then any nonzero vector  $\begin{bmatrix} x \\ x \end{bmatrix}$  is an eigenvector for the eigenvalue 2:

$$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ x \end{bmatrix} = \begin{bmatrix} 2x \\ 2x \end{bmatrix}.$$

Also, any nonzero vector  $\begin{bmatrix} x \\ -x \end{bmatrix}$  is an eigenvector for the eigenvalue 0:

$$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ -x \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Note that an eigenvalue can be 0, but an eigenvector must be nonzero.

## Eigenspaces

If  $\lambda$  is an eigenvalue of the  $n \times n$  matrix  $A$ , we find corresponding eigenvectors by solving the equation

$$(A - \lambda I)\mathbf{x} = \mathbf{0}.$$

The set of *all* solutions is just the null space of the matrix  $A - \lambda I$ .

### Definition

Let  $A$  be an  $n \times n$  matrix, and let  $\lambda$  be an eigenvalue of  $A$ . The collection of all eigenvectors corresponding to  $\lambda$ , together with the zero vector, is called the eigenspace of  $\lambda$  and is denoted by  $E_\lambda$ .

$$E_\lambda = \text{Nul}(A - \lambda I)$$

### Example 3

As before, let  $B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ . In the previous example, we verified that the given vectors were eigenvectors for the eigenvalues 2 and 0.

To find the eigenvectors for 2, solve for the null space of  $B - 2I$ :

$$\text{Nul} \left( \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} - 2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = \text{Nul} \left( \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \right) = \begin{bmatrix} x \\ x \end{bmatrix}.$$

To find the eigenvectors for the eigenvalue 0, solve for the null space of  $B - 0I = B$ .

You can always check if you've correctly identified an eigenvector: simply multiply it by the matrix and make sure you get back a scalar multiple.

## Eigenvalues of triangular matrix

### Theorem

*The eigenvalues of a triangular matrix  $A$  are the entries on the main diagonal.*

*Proof for the  $3 \times 3$  Upper Triangular Case:*

Let

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{33} \\ 0 & 0 & a_{33} \end{bmatrix}.$$

Then

$$A - \lambda I = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{33} \\ 0 & 0 & a_{33} \end{bmatrix} - \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix} = \begin{bmatrix} a_{11} - \lambda & a_{12} & a_{13} \\ 0 & a_{22} - \lambda & a_{23} \\ 0 & 0 & a_{33} - \lambda \end{bmatrix}.$$

By definition,  $\lambda$  is an eigenvalue of  $A$  if and only if  $(A - \lambda I)\mathbf{x} = \mathbf{0}$  has non trivial solutions.

This occurs if and only if  $(A - \lambda I)\mathbf{x} = \mathbf{0}$  has a free variable.

Since

$$A - \lambda I = \begin{bmatrix} a_{11} - \lambda & a_{12} & a_{13} \\ 0 & a_{22} - \lambda & a_{23} \\ 0 & 0 & a_{33} - \lambda \end{bmatrix}$$

$(A - \lambda I)\mathbf{x} = \mathbf{0}$  has a free variable if and only if

$$\lambda = a_{11}, \quad \lambda = a_{22}, \quad \text{or} \quad \lambda = a_{33}$$

An  $n \times n$  matrix  $A$  has eigenvalue  $\lambda$  if and only if the equation

$$A\mathbf{x} = \lambda\mathbf{x}$$

has a nontrivial solution.

Equivalently,  $\lambda$  is an eigenvalue if  $A - \lambda I$  is not invertible.

Thus, an  $n \times n$  matrix  $A$  has eigenvalue  $\lambda = 0$  if and only if the equation

$$A\mathbf{x} = 0\mathbf{x} = \mathbf{0}$$

has a nontrivial solution.

This happens if and only if  $A$  is *not invertible*.

- The scalar 0 is an eigenvalue of  $A$  if and only if  $A$  is *not invertible*.

### Theorem

Let  $A$  be an  $n \times n$  matrix. If  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r$  are eigenvectors that correspond to distinct eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_r$ , then the set  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$  is linearly independent.

The proof of this theorem is in Lay: Theorem 2, Section 5.1.

### Example 4

Consider the matrix

$$A = \begin{bmatrix} 4 & 2 & 3 \\ -1 & 1 & -3 \\ 2 & 4 & 9 \end{bmatrix}.$$

We are given that  $A$  has an eigenvalue  $\lambda = 3$  and we want to find a basis for the eigenspace  $E_3$ .

*Solution* We find the null space of  $A - 3I$ :

$$A - 3I = \begin{bmatrix} 1 & 2 & 3 \\ -1 & -2 & -3 \\ 2 & 4 & 6 \end{bmatrix} \xrightarrow{\text{rref}} \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

$$A - 3I \xrightarrow{\text{rref}} \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

So we get a single equation

$$x + 2y + 3z = 0 \quad \text{or} \quad x = -2y - 3z$$

and the general solution is

$$\mathbf{x} = \begin{bmatrix} -2y - 3z \\ y \\ z \end{bmatrix} = y \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} + z \begin{bmatrix} -3 \\ 0 \\ 1 \end{bmatrix}$$

Hence  $\mathcal{B} = \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -3 \\ 0 \\ 1 \end{bmatrix} \right\}$  is a basis for  $E_3$ .